

Volume 7
Issue 1
October 2022
ISSN:5101195-3

KOSOVA JOURNAL OF SURGERY



- Perspective: RIFAT LATIFI: On Being Minister of Health, Medical Diplomacy and the Transformation of Healthcare in Kosova
- SELMAN URANEUS: Damage Control Surgery in Severe Trauma
- JORG TEBAREK: Aortoiliac treatment with IVUS guidance as a standalone imaging solution: is it feasible and where are the benefits?
- JAMES HU: Familial Adenomatous Polyposis: Review of Current Diagnosis, Screening and Management



KOSOVA JOURNAL OF SURGERY
Volume 7, Issue 1, October 2022
ISSN:5101195-3
info@koscs.org • www.koscs.org

EDITOR-IN- CHIEF

Rifat Latifi, MD, USA

DEPUTY EDITORS-IN-CHIEF

Luan Jaha, MD, Kosova
Nexhmi Hyseni, MD, Kosova
Ramadan Jashari, MD, Belgium
Skender Topi, MD, Albania

ASSISTANT EDITORS

Fatjona Jahaj, Kosova

EDITOR, BIOSTATISTICS

Ilir Hoxha, MD, Kosova

EDITOR OF SURGICAL HISTORY

Isuf Dedushaj, MD, Kosova

HONORARY EDITORS

Sadri Bajraktari, MD, Kosova

DESIGN AND LAYOUT

Ardian Veliu, Kosova

This number is supported by
Swiss Agency for Development and
Cooperation SDC



© Copyright Kosovo Journal of Surgeons.
All rights reserved. No parts of this magazine may be reproduced or transmitted in any form an by any means without written permission from Kosovo Journal of Surgeons. Kosovo Journal of Surgeons reserves the right to reject or accept any advertisement or advertisement material prior to publication.

EDITORIAL BOARD

INTERNATIONAL

Alban Neziri, MD, Switzerland
Alberto R. Ferreres, MD,FACS Argentina
Arben Baboci, MD, Albania
Arben Beqiri, MD, Albania
Arben Gjata, MD, Albania
Artur Xhumani, MD, Albania
Carlos Mesquita, MD, Portugal
Demetrius Litwin, MD, FACS, FICS,USA
Dritan Alushani, MD, Albania
Emir Haxhija, MD, Austria
Fatmir Dalladaku MD, Germany
Georgious Tsoulfas, MD, FACS, FICS, Greece
Gezim Dushaj, MD, Switzerland
Girma Tefera, MD FACS, USA
Giuseppe Nigri, MD, FACS, FRCS, Italy
Hayato Kurihara, MD, FACS, Italy
Horacio Rodriguez Rilo, MD, FACS, USA
Itamar Askenazi, MD, Israel
Joseph Bellal, MD, FACS, USA
Jörg Tessarek, MD, Germany
Kastriot Haxhirexha, MD,North Macedonia
Kita Sallabanda, MD, Spain
Lumnije Kqiku, DDs, Phd, Austria
Maria Castaldi, MD, USA
Minir Asani, MD, Germany
Mustafa Ozsoy, MD, Turkey
Moritz Felsenreich, MD, Austria
Osman Dilek Nuri, MD, Turkey
Peter Killcomons, MD, USA
Philip Caushaj, MD, FACS, USA
Ridvan Alimehmeti, MD, Albania
Russel Andrews, MD, USA
Roberto Bergamaschi, MD, FACS, USA
Salomone Di Saverio MD, FACS, FRCS, England
Selami Sozubir, MD, Turkey
Selman Uranes, MD, FACS, Austria
Tilsed Jonathan, MD, England
Valon Baraliu, MD, Germany
Xheladin Draqini, MD, Albania.

NATIONAL

Agreta Gashi- Gecaj, MD, Kosova
Antigona Hasani, MD, Kosova
Astrit Hamza, MD, Kosova
Avdyl Krasniqi, MD, Kosova
Aziz Mustafa, MD, Kosova
Basri Lenjani, MD, Kosova
Bedri Braha, MD, Kosova
Belinda Pustina, MD, Kosova
Besnik Bicaj, MD, Kosova
Cen Bytyqi, MD, Kosova
Dafina Mahmutaj, MD, Kosova
Dukagjin Spanca, MD, Kosova
Enver Fekaj, MD, Kosova
Eqrem Shala, MD, Kosova
Fadil Beka, MD, Kosova
Fahredin Veselaj, MD, Kosova
Faton Hoxha, MD, Kosova
Fatos Sada, MD, Kosova
Fehmi Ahmeti, MD, Kosova
Ferat Sallahu, MD, Kosova
Gani Çeku, MD, Kosova
Harieta Zherka-Saraqini, MD, Kosova
Hysni Jashari, MD, Kosova
Isa Haxhiu, MD, Kosova
Laura Leci, MD, Kosova
Lulzim Vokrri, MD, Kosova
Nazmi Morina, MD, Kosova
Nehat Baftiu, MD, Kosova
Nexhat Rexha, MD, Kosova
Osman Zhuri, MD, Kosova
Raif Cavolli, MD, Kosova
Sami Salihu, MD, Kosova
Sejdi Statovci, MD, Kosova
Shefqet Lulaj, MD, Kosova
Shqiptar Demaci, MD, Kosova
Sueda Latifi, MD, Kosova
Talat Gjnjolli, MD, Kosova
Vjollca Bince, MD, Kosova
Xhevdet Cuni, MD, Kosova
Xhevdet Tahiraj, MD, Kosova
Ylber Zejnullahu, MD, Kosova

KOSOVA COLLEGE OF SURGEONS

KOSOVA JOURNAL OF SURGERY

October 2022 | Volume 7 | Issue 1 | www.koscs.org

Contents



- 4** Perspective: **Rifat Latifi:** On Being Minister of Health, Medical Diplomacy and the Transformation of Healthcare in Kosova
- 13** **Selman Uraeus:** Damage Control Surgery in Severe Trauma
- 18** **Jorg TeBarek:** Aortoiliac treatment with IVUS guidance as a standalone imaging solution: is it feasible and where are the benefits?
- 25** **Lubomir Martinek:** Ileo-colic Anastomoses – A Review
- 32** **James Hu:** Familial Adenomatous Polyposis: Review of Current Diagnosis, Screening and Management
- 40** **Florin Iordache:** Updates in Fistula-in-Ano. A Narrative Review
- 56** **Selman Uranues, S.** Bile Duct Injury: Prevention Entails Analysis of Why it Happens, Not Only How it Happens?
- 63** **Abbas Smiley:** The Risk of Mortality in Geriatric Patients with Emergent Regional Enteritis is 12-fold Greater than that in Adult Patients: Female Sex, Hospital Length of Stay and Surgery As other Risk Factors of Mortality
- 82** **Adem Miftari:** Intravitreal Anti VEGF Application During Covid – 19 Pandemic in Kosova – Reviewer
- 88** **Abbas Smiley:** Longer stay in hospital is the risk factor for mortality whereas surgery is the protective factor in patients with blood in stool

Updates in Fistula-in-Ano.

A Narrative Review

DR. FLORIN IORDACHE, MD, PHD

Senior Lecturer of General Surgery

University of Medicine and Pharmacy Carol Davila, Bucharest, Romania
Bucharest Emergency Hospital

Abstract

Background: Anal fistula is a challenging disease with a high incidence that affects mainly young people. The aim of this narrative review is to evaluate the recent data regarding the pathogenesis, diagnosis, and treatment of anal fistula patients.

Data sources: The following databases were searched: MEDLINE, Pubmed, Scopus, Cochrane. Relevant articles were selected.

Conclusions: There is a need for better standardization of classifications, measurements, and outcomes. Fistulotomy and fistulectomy are simple and fast with good results in simple fistula. Modern sphincter-sparing techniques are more demanding, and sometimes costly, but there is less risk of incontinence.

Keywords: anal fistula treatment.

Introduction

Anal fistula is defined as a chronic abnormal communication between the epithelialized surface of the anorectal canal and, usually, the perianal skin. Hippocrates made the first reference for fistulous disease, and

he was also the first to advocate the use of a seton (from the Latin *seta*, meaning “bristle”) in about 430 BCE.¹

With a variable annual incidence affecting 8.6-10/100 000 people and a prevalence for whole etiologies (cryptoglandular, Crohn’s etc.) of 1.69 per 10,000 population in the European Union, fistula-in-ano (anal fistula, perianal fistula) is a common cause for surgery.²⁻³ The condition, having a peak between the third to fifth decade of life is prevalent in males.⁴

Carrying the risk of recurrence and fecal incontinence, anal fistula negatively impacts the quality of life of the patient and is a challenge for the surgeon. There is no clear consensus about the optimal surgical procedure. There is also high heterogeneity in outcome selection, definition, and measurement in studies assessing the treatment of cryptoglandular anal fistula.⁵ The aim of this narrative review is to evaluate the recent data in pathogenesis, diagnosis, and treatment of an anal fistula.

Method

A literature search of the last 5 years was performed. The following databases were searched: MEDLINE, Pubmed, Scopus, Cochrane. The search terms included:

“anal fistula,” “anal fistula treatment,” “fistula-in-ano,” “VAAFT,” “LIFT,” and “Laser closure.” Relevant articles were screened, and from a total of 144 titles, the selection included 59 observational studies and 19 randomized controlled trials. Relevant articles prior to the search period were also taken into review.

Pathogenesis

The etiology is not completely understood; however, the most accepted theory is the crypto-glandular theory, initially proposed by Parks.⁴⁻⁵ According to this theory, glandular infections arise from the anal crypts communicating with anal proctodeal glands situated in the intersphincteric space. Dilated glands, which terminate deeper discharge between muscular fibers of the external anal sphincter, or through the external anal sphincter into the ischioanal fossa, will result with anal fistula. This may explain why some of perianal abscesses are associated with anal fistula from the beginning, although anal fistula is considered the chronic phase of intersphincteric anal gland sepsis.⁷ Anorectal infection is responsible for 80-90% of the anal fistula.⁷ Between 30-70% of the patients with anorectal abscesses present concomitantly with a fistula.⁸ More than one third of those who do not present with this condition will be diagnosed with an anal fistula in a period of months or years after abscess drainage.⁸⁻⁹ Less frequently, hemorrhoidectomy, foreign body perforation, and trauma are responsible. Other causes (e.g., Crohn's disease, radiation, tuberculosis etc.) should be thought in suspicious as well when looking at anal fistula, and are at the origin of the remaining 10-20% of the cases.

Some characteristics of anal fistulas cannot be explained by cryptoglandular theory (e.g., many abscesses do not result in a fistula, the prevalence in male patients, the absence of epithelization, and the presence of granulation tissue in most of the fistula tracts).⁷⁻⁸

Classification

The Parks classification is the one mostly utilized.¹²⁻¹³ This system is based on the relationship of the fistula tract to the sphincter, which describes four types of anal fistula with various subtypes. Another two types of classification were added in the current classification (see below):

1. Intersphincteric - the fistula tract ramifies only in the intersphincteric plane; it is the most common of all types, accounting for 55-70% of the cases (45% in the study of Parks).^{1,12,15} This can be further classified as:

- a. Simple intersphincteric, where the fistula tract in the intersphincteric plane passes downward to the anal verge. It is the most frequent subtype of intersphincteric anal fistula; it is considered a low anal fistula.
- b. Intersphincteric fistula with a high blind tract.
- c. Intersphincteric fistula with a high tract opening into the lower rectum.
- d. High intersphincteric fistula without a perineal opening.
- e. High intersphincteric fistula with a pelvic extension.
- f. Fistula from pelvic disease.

2. Transsphincteric – the fistula tract passes from the intersphincteric plane through the external sphincter complex at varying levels into the ischioanal fossa (25-30% of the cases). It can be:

- a. Uncomplicated
- b. With a high blind tract

3. Suprasphincteric - the tract passes in the intersphincteric plane over the top of the puborectalis muscle, then downwards again through the levator plate to the ischioanal fossa, and finally to the skin. Although this type was observed by Parks et al. in 20% of the cases in their series, it is rarely reported today (1-3%).^{1,15} This is further subdivided into:

- a. Uncomplicated
- b. With a high blind tract

4. Extrasphincteric fistulas are very rare (1-5%). The fistula tract passes from the perineal skin through the ischioanal fat and levator muscles into the rectum. It is usually the consequence of trauma or pelvic diseases (e.g., Crohn's). It can be further classified as:

- a. Secondary to transsphincteric fistula
- b. Secondary to trauma
- c. Secondary to anorectal disease (e.g., Crohn's disease)
- d. Secondary to pelvic inflammation

5. Combined

6. Horseshoe – which can occur in up to 4,4% of the cases.¹⁶ It can be:

- a. Intersphincteric
- b. Transsphincteric

Some authors add submucous (subanodermal) fistula as another type of anal fistula. In this case, the tract runs under the anoderm or the rectal mucosa without any involvement of the sphincter.¹⁸ Others consider this type a misnomer.^{1,6}

In Japan, the Sumikoshi classification is more commonly used. In this classification, the space above the dentate line is referred to as high (H); below the dentate line is defined as low (L). Multiple or curved tracts are complex and the letter C is assigned; straight tracts are simple and signified by the letter S. The U letter is used for tracts that extend on one side (unilateral), whereas tracts on both sides are bilateral, and signified with the letter B.¹⁴

Anatomically, anal fistulas can be described as low or high depending on their relationship with the sphincter complex. There is some inconsistency in defining an anal fistula as high. In the classic study of Parks, a high anal fistula passes over the anorectal ring (puborectal muscle). In the consensus statement of the Association of Coloproctologists of Great Britain and Ireland (ACG-BI), a high intersphincteric fistula is defined as a fistula that passes deep to more than 50% of the length of the internal sphincter.⁶ High transsphincteric, suprasphincteric, and extrasphincteric fistulas defined as such pass deep to more than 30% of the external sphincter.^{4,8}

Fistulas that pose a high risk for incontinence are termed complex. A complex fistula-in-ano is defined by one of the following¹⁸:

- when the track crosses, > 30–50% of the external sphincter (high transsphincteric, suprasphincteric or extrasphincteric).
- is anterior in a female.
- is recurrent.
- has multiple tracts.
- the patient has pre-existing incontinence, local irradiation, or Crohn's disease.

Diagnostic

Clinical assessment

Pain, swelling, and discharge are the common complaints of patients with anal fistula. Many patients with fistula have a prior history of an abscess. Sometimes the discharge stops because the fistula opening is temporarily closed with a thin layer of epithelium. The fistula then reopens, with secretion occurring again. In rare cases, anal fistula may be seen in anorectal carcinoma, or an anal or rectal cancer can mimic a fistula¹⁹⁻²²; although there are few cases reported, the examiner should be aware of the possibility.

The patient should be examined on a tilt table in the knee-chest position, or Sims. Visual inspection will generally identify the opening of the fistula. When the external opening is adjacent to the anal margin, it suggests an intersphincteric tract. A more laterally located opening would suggest a transsphincteric fistula. The further the distance of the external opening from the anal margin, the greater the chance of a complicated upward extension. Rectal examination with a bi-digital palpation may determine the primary opening and the fistula tract. Probing should be done gently, never forced, to avoid creating a false tract. Sometimes the angulation or stenosis of the tract precludes passing a probe. If infection/inflammation is present, the maneuver should be done under anesthesia where a therapeutic intervention will be performed (examination under anesthesia – EUA).

In general, the internal opening will be found at anoscopy at the level of the dentate line. Gentle pressure on the fistula track may result in discharge of pus from the primary opening. As Goodsall's rule states, when the external opening lies anterior to the transverse plane, the internal opening is located radially most of the time. When the external opening is posterior to the plane, the internal opening is generally (not always) in the posterior midline. A recent prospective study with a total of 188 patients found a predictive accuracy of 84.6% in case of a fistula with an anterior external opening.²³ In those with a posterior external opening, this was found to be 69.1% accurate with an overall predictive accuracy of Goodsall's rule of 77%.²³ As demonstrated, this rule should be used as guide to help the surgeon to find the tract and it is not absolute.^{1,23}

Imaging

Most of the time, clinical examination is sufficient in primary anal fistula seen for the first time. Nevertheless, in complex fistulas, recurrent ones, or after multiple perianal procedures, imaging is necessary. From a practical standpoint, an imaging method should help in classifying the fistula, visualizing the tract and its secondary extensions, the relationship with the sphincter apparatus (its involvement), and to identify the internal opening. Historically, the following methods were used to attain the stated objectives:

- Fistulography – is a radiological study visualizing the fistula tract by injecting a few ml of a hydro-soluble contrast agent. It is seldom used today as it

does not give clear tridimensional anatomical information about the fistula tract in relationship with anatomical structures, mainly the sphincter. This method has a 10% rate of false-positive results, and an accuracy of only 16% in identifying the internal openings and secondary extensions.²⁴

- Computed tomography (CT) - with administration of contrast intravenously, intrarectally, orally, or by injection through the fistula tract is helpful in high fistula (pelvic or supralelevatorian) is superseded today by MRI or endoanal ultrasonography. With an accuracy rate between 24–60% and the burden of radiation, CT is rarely used today.²⁴ CT with intravenous contrast is sometimes still performed where MRI and/or anal US is unavailable, or in an emergency setting.²⁵ In these acute situations, CT-scan may provide useful information about a pelvic infection.
- Ultrasonography - endo-anal ultrasonography (EUS) in anal fistulas was described first by Bartram et al. and the accuracy of the early studies was between 63-87%.^{6,26-27} Using 3-D mode further improved the accuracy to 91-96% in experienced hands.²⁸⁻²⁹ 3D-EUS is a reliable technique usually performed in left lateral decubitus position, although prone position or lithotomy position can also be used.³⁰⁻³¹ It can accurately classify fistulas and report the primary tract, the location of the internal opening, and the relation of the fistula to the anal sphincter muscles. In a prospective study of 196 cases with anal fistula comparing the clinical and functional outcome (incontinence), the usefulness of 3D-EUS was validated.²⁸ With a follow-up of up to one year in the two matched subgroups, one group with 3D-EUS and the other with clinical examination under anesthesia, Ding et al. demonstrated a statistical reduction of the recurrence rate in the EUS subgroup (8.8% vs. 13.8%).²⁸ The difference was even more significant in patients with complex fistulas (12.8% vs. 22.5%).³² Additionally, a significant decrease in continence was found after one year in patients that were not evaluated with EUS.³² Due to the high frequency used for EUS (between 7-10 MHz), the loss of resolution is important farther away from the transducer; this limits the value of EUS in visualizing secondary extensions, the ischioanal and ischiorectal region, and the supralelevator area. Also, because of fibrosis and

sclerosis encountered in recurrent fistulae, EUS is of limited help. Using hydrogen peroxide or a contrast agent injected through the fistula opening can enhance imaging in EUS thus better displaying the fistula tract.^{25,27} In a very recent study, ultrasound with contrast media producing acoustic scattering of microbubble to enhance contrast was compared with standard MRI in 98 patients.³³ Both methods exhibited similar accuracy, but the accuracy of contrast perineal ultrasound was better than MRI for identifying an internal opening under 3 cm from the anal verge.³³ Still, hydrogen peroxide has the advantages of availability and cost as compared with any other contrast media. As an alternative to EUS, transperineal ultrasonography (TPUS) showed comparable, or even better than, EUS results in defining and classifying anal fistulae.³⁴ In an early study comparing EUS and TPUS in patients with perianal Crohn's disease, with EUS as the reference standard, TPUS had a sensitivity of 84.9% and a positive predictive value of 86.5% for anal fistulas and was of similar value as EUS for the diagnosis of anorectal abscess.³⁵ In another recent systematic review with a meta-analysis evaluating transperineal ultrasound (TPUS), the pooled results for perianal fistula identification showed a sensitivity of 98% and a positive predictive value (PPV) of 95% (95% CI 83–98%).³⁶ In the same review, the usefulness of TPUS for perianal fistulae classification has been found with a pooled sensitivity and a PPV of 92%.³⁶ The sensitivity for the classification of fistulae was higher when the comparator was examined under anesthesia (98%) rather than transrectal ultrasound or MRI (87%). A drawback of this review was the significant heterogeneity among the six studies included in the review.³⁶ TPUS has several important advantages in the diagnosis and classification of anal fistula, such as: relative ease of use, minimal discomfort, cost-effectiveness, relatively non-time consuming, and widely available in the clinical setting. Although recent data are encouraging, there is still no consensus about its use in perianal fistula. A common limitation for both EUS and TPUS is that they are both operator dependent.

- Magnetic resonance imaging - it is considered the “golden standard,” and has the advantage of not being user-dependent for interpretation, as well as the ability to evaluate fistula tracts that course distant

(less than 4 weeks); and an associated stricture of the anorectum are relative contraindications to the rectal advancement flap.⁴⁷ Different forms of flaps (rhomboids, elliptical) are utilized by different authors without evidence of better results.⁵³ There is data showing that increasing the thickness of the flap improves the healing rates. The healing rate of endoanal advancement flap ranges between 68-87%.⁵⁴ In cases of failure, the procedure can be repeated.^{7,47,55} Interestingly, patients with anal fistulas with and without Crohn's disease exhibit the same healing rate.⁵⁶

Video-assisted anal fistula treatment (VAAFT)

Video-assisted anal fistula treatment is one of the sphincter-sparing procedures. VAAFT was described by Meinero and Mori in 2011, and has a diagnostic phase and a treatment phase.⁵⁷ Closing the internal opening is essential for success; it can be performed with a stapler, suture, or a flap (mucous or cutaneous). Initial results in 136 patients with anal fistula showed primary healing in 73 patients (73.5%) in the following 3 months after the procedure. No healing was found in 26 patients (26.5%).⁵⁷ Further reinterventions were executed and VAAFT was implemented again. At 12 months, the authors reported a healing rate of 87.1%.⁵⁷ Compared with FiLaC (Fistula Laser Closure (FiLaC™)), the advantage of the procedure is that it is performed under direct vision.

During the reviewed period, 9 observational studies were found to meet the search criteria. Data from these studies is presented in tables 2 and 3. The healing rate of VAAFT was between 51-89.5%, and the recurrence rate was between 11.5 - 35% (tables 2 and 3). There were differences among the studies regarding the operative phase of the procedure, particularly the treatment of the internal opening (table 2). While heterogeneity of the studies is evident, the pooled healing rate was only 81.2%, and the reported rate of complications was 10.7%. Most complications were grade 1 on the Clavien-Dindo scale, although there were also grade 2 and even grade III type complications reported (table 3). Scrotal/perineal oedema or bleeding were the most severe reported complications.

In a very recent randomized control trial, Sorensen et al. compared VAAFT with fistulectomy and sphincter repair (FSR) in terms of wound healing, functional outcome, and quality of life.⁵⁸ Due to the high recurrence in both groups (65% for VAAFT and 27% for FSR), the study was terminated early.⁵⁸ It would seem that VAAFT

fails to increase the healing rate in complex anal fistulas. It is also worth noting that in cases of VAAFT, there is a minimal risk of incontinence.

Fistula Laser Closure (FiLaC™)

FiLaC as a novel sphincter-preserving technique closure of the fistula tract using laser ablation was first published in 2011 by Wilhelm.⁵⁹ In this first study, healing occurred in 9 of the 11 patients involved the study (81.6%).⁵⁹ In the FiLaC procedure, the fistula tract is destroyed using a radial-emitting laser fiber. It can be combined with an additional method to close the internal orifice. Laser fistulectomy has increased in popularity due to its minimal safety issues, limited side effect profile, and short learning curve; however, the results are still suboptimal in some studies (see table 4).

Ten studies were included in the present review (table 4). Overall, 623 patients were included, from which only 38 had Crohn's disease (6%) and 468 (74.5%) had previous surgery. The extremely variable primary healing rate was between 22-74.7% (table 4). The overall healing rate attained a maximum of 88% in the study of Wilhelm et al. with a follow-up of five years.⁶⁰ In the same study, a total rate of complication of 7.7% was also noted. Similar to other studies analyzing anal fistula treatment, the heterogeneity and the selection of the cases made interpretation of the results difficult. Moreover, there were differences in techniques among different groups in terms of power delivered (between 10-14 W), the speed of retracting the fiber, and the usage of different types of probes. Some authors consider that healing rates are better when the internal opening is closed, which tends to be the norm.⁶⁰ Others refute this theory.⁶¹ It may well be that the reason for such variety in the healing rates is the length of the fistula tract, as demonstrated by Lauretta et al.⁶² In their retrospective observational study of 30 patients using a length of 30 mm as a cutoff, the shorter fistulas had a significantly better prognosis: 58.3% (7 out of 12) for fistulas shorter than 30 mm and 16.6% (3 out of 18) for fistulas longer than 30 mm.⁶² In a recent systematic review and meta-analysis, the pooled success rate of the technique was 63%, and the pooled complication rate was 8%.⁶³ It can be concluded that FiLaC™ has success rates comparable to other sphincter-sparing techniques. The technique is safe with respect to adverse events and risk of incontinence, though it has no significant advantage over other methods that have been demonstrated up until now.

Ligation of the intersphincteric fistula tract (LIFT)

The LIFT technique starts with an incision along the intersphincteric groove, and meticulous dissection is performed to identify the fistula tract. This is ligated, and the fistula and the tissue of the fistula wall is cut and cleared in this way in order to eliminate the septic focus. It is considered appropriate for transsphincteric fistulas, including most complex anal fistulas, recurrent, and fistulas that fail after other surgical procedures.⁵¹ Rojanasakul et al. introduced the LIFT technique as a cost-effective sphincter sparing method in 2007.⁶⁴ In a more recent study from the same center assessing the experience with LIFT over 10 years, the healing rate was 87.7%.⁶⁴ In a recent systematic review and meta-analysis, Emile et al. focused on a total of 1376 patients with a follow-up of one year and found an overall success rate of 78% (1075 patients) for LIFT and an incontinence rate of 1.4%, all but minor.⁶⁵ Horseshoe fistula, fistulas associated with Crohn's disease, and a history of previous fistula surgery were associated with a greater incidence of persistence or recurrence after LIFT.⁶⁵ It must be pointed out that, in the same study, the weighted mean complication rate was 13.9% higher than previously thought. The most common complications after LIFT were dehiscence and infection.⁶⁵ Leaving the wound open may avoid these complications. Based on their data, the conclusion of the systematic review concluded that LIFT had a better success rate and less complications than the rectal advancement flap technique.⁶⁶ In the recent randomized controlled trials presented in this review, the healing rate of LIFT was between 77.5-93.3% (table 5). Different modifications of the LIFT procedure were also proposed. Chen et al. used a lateral incision to access the intersphincteric groove in 10 patients.⁶⁷ In this modification, after dissection, the proximal end of the fistula was clamped above the external sphincter, and the distal portion was removed for pathological examination. The stump is ligated with a resorbable thread. The advantage of this approach is is a better view of the dissection.⁶⁷ Using the same approach, Kang et al. attained a healing rate of 75% (21 patients from 28).⁶⁸ One other modification of the LIFT procedure consisted of associating an advancement flap to the LIFT. The only report addressing this technique found a lower healing rate, primary or secondary to the original LIFT procedure.⁶⁹ Another modification focused on associating a bioprosthetic (resorbable) material in the intersphincteric plane to buttress the closure of the intersphincteric tract, also known as Bio-LIFT, originally

described by Ellis et al.⁷⁰ Nonetheless, this procedure is expensive and requires a more extensive dissection in the intersphincteric space than is usual for the LIFT. Another modification of LIFT plus plug was performed by Han et al. and involved the excision of a section of fistula tract in the intersphincteric space.⁷¹ Then, the bioprosthetic plug (porcine small-intestine submucosa extracellular matrix plug) is placed into the fistula tract through an opening in the external sphincter to the external opening in the skin. In their randomized controlled trial, the rate of success for the modified procedure was 94% compared with 83.9% for simple LIFT.⁷¹ In another retrospective cohort comparing BioLIFT with the LIFT procedure, Zwiép et al. found that the primary healing rate was 75% versus 58.7%, and the complication rate was 22.7% versus 17.3% in favor of the former.⁷² Madbouly et al. recently combined injecting autologous platelet plasma rich with the LIFT procedure.⁷³ Their results showed a better healing rate for the modified LIFT procedure (77.5% vs. 59% - table 5). The LIFT procedure remains an adequate option as a sphincter-sparing procedure in anal fistulas. One advantage of LIFT could be understood as the downstaging of the fistula from transsphincteric to intersphincteric, which seems to take place in a proportion of failures.⁴⁷ This allows for a simpler treatment, thus preserving the external sphincter which would have been involved originally.

Anal fistula plug (AFP)

Another sphincter-sparing technique is an anal fistula plug. A variety of materials, bio-prosthetic or synthetic materials, are used to occlude the fistula track (human acellular dermal matrix, porcine dermal collagen etc.), by blocking the internal opening.⁷⁴ In this way, fecal material cannot enter the fistula tract to promote healing. Existing data do not demonstrate a better healing rate for this procedure. In a randomized controlled trial accruing a total of 304 patients, Jayne et al. found clinical healing rates associated with the fistula plug and surgeon's preference groups to be 54% and 55% at the 12-month follow-up, respectively.⁷⁵ Additionally, there were higher costs and highly uncertain quality-adjusted life-year (QALY) gains associated with the fistula plug.⁷⁵ In a systematic review comparing the rectal advancement flap and anal fistula plug procedure after pooling RCTs with long term follow-up, found that in the treatment of complex cryptoglandular anal fistulas, the RAF was superior to AFP in terms of healing and

recurrence rate.⁷⁶ Plug dislodgement or sepsis are the most common causes of failure. A variant of the plug for anal fistula is filling the fistula tract with material that will block communication of the fistula tract with the anorectum and promote healing. Fibrin glue was the first material to be used. The healing rates with fibrin glue are extremely variable, ranging between 14% to 94%.⁴⁷ Other filling materials were tried, including platelet-rich plasma, autologous cartilage, fat, autologous micro-fragmented adipose tissue, alginate, and allogeneic bone marrow-derived mesenchymal stromal cell.^{51,77} There are high rates of recurrence with filling therapy, although there are differences among the different types of materials used. There is no strong evidence supporting these methods.

OTSC®(Over-the-Scope-Clip) Proctology Device

The OTSC™ Proctology system (Ovesco Endoscopy AG, Tuebingen, Germany) consists of a clip and applicator, and is a modification of the endoscopic OTSC™ clip, which is made of a super elastic shape memory alloy (Nitinol). When the opened clip is released from the applicator tip, it automatically closes and exerts constant pressure on the tissue between the jaws of the clip. When applied on the internal fistula opening, the clip compresses and seals the orifice. In this way, the fecal content from the rectum is blocked from entering the fistula tract maintaining the sepsis. It is considered a sphincter-sparing minimally invasive procedure. The possible migration of the clip and postoperative abscesses are the specific possible complications for this procedure.⁷⁸ In one of the first studies of this method reported that the healing rate was only 18%.⁷⁸ In another more recent study, the OTSC™ Proctology system attained a healing rate of 80%.⁷⁹ In the only review of this method (over 200 cases), the healing rate was 63%.⁸⁰ Besides being a minimally invasive procedure and a sphincter sparing one, closing the fistula with this special clip has the advantages of repeatability as well as a short operative time; however, further studies are needed to determine the advantages.⁸⁰

New procedures

New surgical treatments for anal fistulas were proposed by Garg et al.^{15,81} The first procedure consisted of proximal superficial cauterization, regularly emptying fistula tracts and curettage of tracts (PERFACT, as was named by the author) procedure, and initially displayed a healing rate of 86%, which later dropped to 50%.¹⁵

Subsequently, the same authors proposed transanal opening of intersphincteric space (TROPIS) for high complex fistulas.¹⁵ Transanally, the internal opening and the intersphincteric portion of the fistula tract were both laid open into the anal canal. The resulting wound was left open to heal by secondary intention. The external tracts were curetted, and a drainage tube was placed in the cleaned tracts from the external opening up to the lateral border of the external sphincter. Healing occurred in 86% and recurrence in 14%.¹⁵ The data was determined from a single center TROPIS procedure and was not replicated by others until now.

Conclusion

Further standardization of definitions, classifications, measurements, and outcomes in anal fistula are needed. Tailoring the treatment is still the best strategy to get optimal results. Fis tulotomy and fistulectomy are simple, fast, and provide adequate results in simple fistula, but with a higher risk of incontinence. Associating marsupialization improves the healing process. Sphincter-sparing techniques can be more technically demanding, and therefore operating time can be longer, but with less risk of incontinence. Despite technological advances and increasing cost of treatments, the search for the ideal treatment of anal fistula continues.

Conflict of Interest Disclosure Statement

The author reports no disclosures.

REFERENCES

1. Ky AJ, Steinhagen E. Anal Fistula. In: M Corman, R Bergamaschi, J Nicholls, V Fazio E, editor. Corman's Colon and Rectal Surgery. Sixth Edit. Philadelphia, PA: Lip-pincott Williams & Wilkins; 2012. p. 384-427.
2. Kontovounisios C, Tekkis P, Tan E, Rasheed S, Darzi A, Wexner SD. Adoption and success rates of perineal procedures for fistula-in-ano: A systematic review. *Colorectal Disease*. 2016;18(5):441-58.
3. García-Olmo D, Assche G van, Tagarro I, et. al. Prevalence of Anal Fistulas in Europe: Systematic Literature Reviews and Population- Based Database Analysis. *Advances in Therapy [Internet]*. 2019 Dec 1 [cited 2021 Sep 11];36(12):3503. Available from: /pmc/articles/PMC6860471/
4. Hokkanen SR, Boxall N, Khalid JM, Bennett D, Patel H. Prevalence of anal fistula in the United Kingdom. *World Journal of Clinical Cases [Internet]*. 2019 Jul 1 [cited 2021 Sep 18];7(14):1795. Available from: /pmc/articles/PMC6692258/

5. Machielsen AJHM, Iqbal N, Kimman ML, et al. Heterogeneity in outcome selection, definition and measurement in studies assessing the treatment of cryptoglandular anal fistula: findings from a systematic review [Internet]. Vol. 25, *Techniques in Coloproctology*. Springer International Publishing; 2021. 761–830. Available from: <https://doi.org/10.1007/s10151-021-02452-5>
6. Williams JG, Farrands PA, Williams AB, et al. The treatment of anal fistula: ACPGBI position statement. *Colorectal Disease*. 2007 Oct;9(SUPPL.4):18–50.
7. Malik AI, Nelson RL, Tou S. Incision and drainage of perianal abscess with or without treatment of anal fistula. *Cochrane Database of Systematic Reviews*. 2010;(7).
8. Vogel JD, Johnson EK, Morris AM, et al. Clinical Practice Guideline for the Management of Anorectal Abscess, Fistula-in-Ano, and Rectovaginal Fistula. *Diseases of the Colon and Rectum*. 2016;59(12):1117–33.
9. Abcarian H. Anorectal Infection: Abscess–Fistula. *Clinics in Colon and Rectal Surgery [Internet]*. 2011 [cited 2021 Sep 26];24(1):14. Available from: [/pmc/articles/PMC3140329/](https://pubmed.ncbi.nlm.nih.gov/2140329/)
10. Gosselink MP, Onkelen RS van, Schouten WR. The cryptoglandular theory revisited. *Colorectal Disease [Internet]*. 2015 Dec 1 [cited 2021 Sep 12];17(12):1041–3. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/codi.13161>
11. Mitalas LE, van Onkelen RS, Monkhorst K, et al. Identification of epithelialization in high transsphincteric fistulas. *Techniques in Coloproctology*. 2012;16(2):113–7.
12. Parks AG, Gordon PH, Hardcastle JD. A classification of fistula-in-ano. *British Journal of Surgery*. 1976;63(1):1–12.
13. de Groof EJ, Cabral VN, Buskens CJ, et al. Systematic review of evidence and consensus on perianal fistula: An analysis of national and international guidelines. *Colorectal Disease*. 2016;18(4):O119–34.
14. Tetsuo Y. Japanese Practice Guidelines for Anal Disorders. *Journal of the Anus, Rectum and Colon*. 2018;2:103–9.
15. Garg P, Kaur B, Goyal A, Yagnik VD, Dawka S, Menon GR. Lessons learned from an audit of 1250 anal fistula patients operated at a single center: A retrospective review. *World Journal of Gastrointestinal Surgery*. 2021;13(4):340–54.
16. Parades V de, Fathallah N, Blanchard P, Zeitoun J-D, Benadji B, Atienza P. Horseshoe tract of anal fistula: bad luck or an avoidable extension? Lessons from 82 cases. *Colorectal Disease [Internet]*. 2012 Dec 1 [cited 2021 Sep 1];14(12):1512–5. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1463-1318.2012.03034.x>
17. Ommer A, Herold A, Berg E, et al. German S3 guidelines: anal abscess and fistula (second revised version). *Langenbeck's Archives of Surgery*. 2017;402(2):191–201.
18. Whiteford MH, Kilkenny J, Hyman N, et al. Practice parameters for the treatment of perianal abscess and fistula-in-ano (revised). *Diseases of the Colon and Rectum*. 2005;48:1337–42.
19. Gaertner WB, Hagerman GF, Finne CO, et al. Fistula-associated anal adenocarcinoma: Good results with aggressive therapy. *Diseases of the Colon and Rectum*. 2008 Jul;51(7):1061–7.
20. Ohta R, Sekikawa K, Goto M, et al. A case of perianal mucinous adenocarcinoma arising from an anorectal fistula successfully resected after preoperative radiotherapy. *Case Reports in Gastroenterology*. 2013 May;7(2):219–23.
21. Montero-Mateos E, Carmen S del, Sanz J, et al. Endoluminal tumor implant of a colorectal cancer in an anal fistula detected by FISH techniques: a case report. *Journal of Gastrointestinal Oncology [Internet]*. 2021 Apr 1 [cited 2021 Sep 1];12(2):900. Available from: [/pmc/articles/PMC8107577/](https://pubmed.ncbi.nlm.nih.gov/38107577/)
22. Zhu X, Zhu TS, Ye DD, Liu SW. Magnetic resonance imaging findings of carcinoma arising from anal fistula: A retrospective study in a single institution. *World Journal of Clinical Cases*. 2020;8(21):5159–71.
23. Devi VS, Thulasibai SKL, Deepak P, Babu A, Jayasankar, Beena K. Goodsall's rule— its predictive accuracy in tracing the tract of fistula in ano. *International Surgery Journal*. 2020 Nov 27;7(12):4116.
24. Yassin NA, Day N, Phillips RKS. Imaging of anal fistulas. *Seminars in Colon and Rectal Surgery [Internet]*. 2014 [cited 2021 Oct 11];25:176–82. Available from: <http://dx.doi.org/10.1053/j.scrs.2014.08.004>
25. Halligan S, Tolan D, Amitai MM, et al. ESGAR consensus statement on the imaging of fistula-in-ano and other causes of anal sepsis. *European Radiology* 2020 30:9 [Internet]. 2020 Apr 19 [cited 2021 Sep 11];30(9):4734–40. Available from: <https://link.springer.com/article/10.1007/s00330-020-06826-5>
26. Kim Y, Park YJ. Three-dimensional endoanal ultrasonographic assessment of an anal fistula with and without H2O2 enhancement. *World Journal of Gastroenterology : WJG [Internet]*. 2009 [cited 2021 Oct 11];15(38):4810. Available from: [/pmc/articles/PMC2761560/](https://pubmed.ncbi.nlm.nih.gov/192761560/)
27. Cao F, Liu G, Wen Y, et al. Application of Endorectal Ultrasound in Evaluation of Anorectal Diseases. *International Journal of Radiology [Internet]*. 2016 Jun 27 [cited 2021 Oct 14];3(2):109–19. Available from: <http://www.ghrnet.org/index.php/IJR/article/view/1609/2071>
28. Kołodziejczak M, Santoro GA, Obcowska A, Lorenc Z, Mańczak M, Sudol-Szopińska I. Three-dimensional endoanal ultrasound is accurate and reproducible in determining type and height of anal fistulas. *Colorectal Disease [Internet]*. 2017 Apr 1 [cited 2021 Oct 20];19(4):378–84. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/codi.13580>

29. Rahmanian E, Frootan M, Anaraki F, Alimadadi E, Ramezanzpoor M. Evaluation of diagnostic accuracy of 3-D endoanal ultrasound in perianal fistula in comparison with intra-operative findings. *Journal of Coloproctology [Internet]*. 2018 Jan 1 [cited 2021 Oct 20];38(1):9–12. Available from: <http://jcol.elsevier.es/en-evaluation-diagnostic-accuracy-3-d-endoanal-articulo-S2237936317300734>
30. Lengyel AJ, Hurst NG, Williams JG. Pre-operative assessment of anal fistulas using endoanal ultrasound. 2002;436–40.
31. Sharma A, Yadav P, Sahu M, Verma A. Current imaging techniques for evaluation of fistula in ano: a review. *Egyptian Journal of Radiology and Nuclear Medicine* 2020 51:1 [Internet]. 2020 Jul 22 [cited 2021 Oct 11];51(1):1–18. Available from: <https://ejrnm.springeropen.com/articles/10.1186/s43055-020-00252-9>
32. Ding JH, Bi LX, Zhao K, et al. Impact of three-dimensional endoanal ultrasound on the outcome of anal fistula surgery: A prospective cohort study. *Colorectal Disease*. 2015;17(12):1104–12.
33. Lin T, Ye Z, Hu J, Yin H. A comparison of trans-fistula contrast-enhanced endoanal ultrasound and MRI in the diagnosis of anal fistula. *Annals of Palliative Medicine [Internet]*. 2021 Aug 1 [cited 2021 Oct 11];10(8):9165173–9173. Available from: <https://apm.amegroups.com/article/view/76216/html>
34. Zbar AP, Oyetunji RO, Gill R. Transperineal versus hydrogen peroxide-enhanced endoanal ultrasonography in never operated and recurrent cryptogenic fistula-in-ano: A pilot study. *Techniques in Coloproctology*. 2006;10(4):297–302.
35. Maconi G, Ardizzone S, Greco S, Radice E, Bezzio C, Bianchi Porro G. Transperineal ultrasound in the detection of perianal and rectovaginal fistulae in Crohn's disease. *American Journal of Gastroenterology*. 2007;102(10):2214–9.
36. Maconi G, Greco MT, Asthana AK. Transperineal Ultrasound for Perianal Fistulas and Abscesses - A Systematic Review and Meta-Analysis. Vol. 38, *Ultraschall in der Medizin*. Georg Thieme Verlag; 2017. p. 265–72.
37. Dekker L, Zimmerman DDE, Smeenk RM, Schouten R, Han-Geurts IJM. Management of cryptoglandular fistula-in-ano among gastrointestinal surgeons in the Netherlands. *Techniques in Coloproctology [Internet]*. 2021;25(6):709–19. Available from: <https://doi.org/10.1007/s10151-021-02446-3>
38. Koelbel G, Schmiedl U, Majer MC, et al. Diagnosis of fistulae and sinus tracts in patients with Crohn disease: Value of MR imaging. *American Journal of Roentgenology*. 1989;152(5):999–1003.
39. Luniss PJ, Armstrong P, Barker PG, Reznick TH, Phillips RKS. Magnetic resonance imaging of anal fistulae. *The Lancet*. 1992;340:394–6.
40. Sudół-Szopińska I, Santoro GA, Kołodziejczak M, Wiaczek A, Grossi U. Magnetic resonance imaging template to standardize reporting of anal fistulas. *Techniques in Coloproctology [Internet]*. 2021;25(3):333–7. Available from: <https://doi.org/10.1007/s10151-020-02384-6>
41. Kumar N, Agarwal Y, Chawla AS, Jain R, Thukral BB. MRI of perianal fistulae: A pictorial kaleidoscope. *Clinical Radiology*. 2015;70(12):1451–61.
42. Morris J, Spencer JA, Ambrose NS. MR Imaging Classification of Perianal Fistulas and Its Implications for Patient Management. <https://doi.org/10.1148/radiographics.203.g00mc15623> [Internet]. 2000 May 1 [cited 2021 Oct 6];20(3):623–35. Available from: <https://pubs.rsna.org/doi/abs/10.1148/radiographics.203.g00mc15623>
43. Konan A, Onur MR, Özmen MN. The contribution of pre-operative MRI to the surgical management of anal fistulas. *Diagnostic and Interventional Radiology*. 2018;24(6):321–7.
44. Garg P, Yagnik VD, Kaur B, Menon GR, Dawka S. Role of MRI to confirm healing in complex high cryptoglandular anal fistulas: long-term follow-up of 151 cases. *Colorectal Disease*. 2021;(March):1–9.
45. Garg P. Comparison of Preoperative and Postoperative MRI After Fistula-in-Ano Surgery: Lessons Learnt from An Audit of 1323 MRI At a Single Centre. *World Journal of Surgery [Internet]*. 2019;43(6):1612–22. Available from: <https://doi.org/10.1007/s00268-019-04926-y>
46. Brilliantino A, Iacobellis F, Reginelli A, et al. Pre-operative assessment of simple and complex anorectal fistulas: Tridimensional endoanal ultrasound? Magnetic resonance? Both? *Radiologia Medica [Internet]*. 2019;124(5):339–49. Available from: <https://doi.org/10.1007/s11547-018-0975-3>
47. Williams G, Williams A, Tozer P, et al. The treatment of anal fistula: second ACPGBI Position Statement – 2018. *Colorectal Disease*. 2018;20:5–31.
48. Anan M, Emile SH, Elgendy H et al. Fistulotomy with or without marsupialisation of wound edges in treatment of simple anal fistula: a randomised controlled trial. [cited 2021 Oct 24]; Available from: www.clinicaltrials.gov
49. Sahebally SM, O'Byrne L, Troy A, Byrnes KG, Burke J, McNamara D. A meta-analysis of marsupialisation versus none in the treatment of simple fistula-in-ano. *International Journal of Colorectal Disease*. 2021;36(3):429–36.
50. Seyfried S, Bussen D, Joos A, Galata C, Weiss C, Herold A. Fistulectomy with primary sphincter reconstruction. *International Journal of Colorectal Disease*. 2018 Jul 1;33(7):911–8.
51. Ji L, Zhang Y, Xu L, Wei J, Weng L, Jiang J. Advances in the Treatment of Anal Fistula: A Mini-Review of Recent Five-Year Clinical Studies. *Frontiers in Surgery*. 2021 Feb 11;7 (February):1–9.
52. Elting AW. The treatment of fistula in ano. *Annals of surgery*. 1912;56(5):744–52.

53. Yellinek S, Krizzuk D, Moreno Djadou T, Lavy D, Wexner SD. Endorectal advancement flap for complex anal fistula: does flap configuration matter? *Colorectal Disease*. 2019;21(5):581–7.
54. Zimmerman DDE, Stijns J, Wasowicz DK, Gottgens KWA. Transanal Advancement Flap Repair: The Current Gold Standard for Cryptoglandular Transsphincteric Perianal Fistulas. *Turkish Journal of Colorectal Disease*. 2019;29(3):104–10.
55. Stremitzer S, Riss S, Swoboda P, Pozios I. Repeat endorectal advancement flap after flap breakdown and recurrence of fistula-in-ano – is it an option? *Colorectal Disease [Internet]*. 2012 Nov 1 [cited 2021 Sep 1];14(11):1389–93. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1463-1318.2012.02990.x>
56. Seifarth C, Lehmann KS, Holmer C, Pozios I. Healing of rectal advancement flaps for anal fistulas in patients with and without Crohn’s disease: a retrospective cohort analysis. *BMC Surgery [Internet]*. 2021;21(1):1–7. Available from: <https://doi.org/10.1186/s12893-021-01282-4>
57. Meinerio P, Mori L. Video-assisted anal fistula treatment (VAAFT): a novel sphincter-saving procedure for treating complex anal fistulas.
58. Sørensen, KM, Möller S, Qvist N, et al. Video-assisted anal fistula treatment versus fistulectomy and sphincter repair in the treatment of high cryptoglandular anal fistula: a randomized clinical study. *BJS Open [Internet]*. 2021 Sep 6 [cited 2021 Oct 26];5(5):1–7. Available from: <https://academic.oup.com/bjsopen/article/5/5/zrab097/6382011>
59. Wilhelm A. A new technique for sphincter-preserving anal fistula repair using a novel radial emitting laser probe. *Techniques in Coloproctology*. 2011;15(4):445–9.
60. Wilhelm A, Fiebig A, Krawczak M. Five years of experience with the FiLaC™ laser for fistula-in-ano management: long-term follow-up from a single institution. *Techniques in Coloproctology*. 2017;21(4):269–76.
61. Serin KR, Hacim NA, Karabay O, Terzi MC. Retrospective Analysis of Primary Suturing of the Internal Orifice of Perianal Fistula during FiLaC Procedure. *Surgical Laparoscopy, Endoscopy and Percutaneous Techniques*. 2020;30(3):266–9.
62. Lauretta A, Falco N, Stocco E, Bellomo R, Infantino A. Anal Fistula Laser Closure: the length of fistula is the Achilles’ heel. *Techniques in Coloproctology [Internet]*. 2018;22(12):933–9. Available from: <http://dx.doi.org/10.1007/s10151-018-1885-z>
63. Frountzas M, Stergios K, Nikolaou C, et al. Could FiLaC™ be effective in the treatment of anal fistulas? A systematic review of observational studies and proportional meta-analysis. *Colorectal Disease*. 2020;22(12):1874–84.
64. Malakorn S, Sammour T, Khomvilai S, et al. Ligation of intersphincteric fistula tract for fistula in Ano: Lessons learned from a decade of experience. *Diseases of the Colon and Rectum*. 2017 Oct 1;60(10):1065–70.
65. Emile SH, Khan SM, Adejumo A, Koroye O. Ligation of intersphincteric fistula tract (LIFT) in treatment of anal fistula: An updated systematic review, meta-analysis, and meta-regression of the predictors of failure. *Surgery (United States) [Internet]*. 2020;167(2):484–92. Available from: <https://doi.org/10.1016/j.surg.2019.09.012>
66. Hany Emile S, Elfeki H, Shalaby M, Sakr A. A Systematic review and meta-analysis of the efficacy and safety of video-assisted anal fistula treatment (VAAFT). *Surg Endosc*. 2018;32:2084–93.
67. Chen T-A, Liu K-Y, Yeh C-Y. High ligation of the fistula track by lateral approach: a modified sphincter-saving technique for advanced anal fistulas. *Colorectal Disease [Internet]*. 2012 Sep 1 [cited 2021 Sep 1];14(9):e627–30. Available from: <https://www.onlinelibrary.wiley.com/doi/full/10.1111/j.1463-1318.2012.03050.x>
68. Kang WH, Yang HK, Chang HJ, et al. High ligation of the anal fistula tract by lateral approach: A prospective cohort study on a modification of the ligation of the intersphincteric fistula tract (LIFT) technique. *International Journal of Surgery*. 2018 Dec 1;60:9–14.
69. Hussain T, Hughes B, Sharma A. LIFT combined with mucosal advancement flap for transsphincteric fistulas. *Techniques in Coloproctology [Internet]*. 2019;23(3):289–90. Available from: <http://dx.doi.org/10.1007/s10151-019-01956-5>
70. Ellis CN. Outcomes with the use of bioprosthetic grafts to reinforce the ligation of the intersphincteric fistula tract (BioLIFT procedure) for the management of complex anal fistulas. *Diseases of the Colon and Rectum*. 2010;53(10):1361–4.
71. Han G, Zhen JW, Yi Z, et al. Ligation of intersphincteric fistula tract vs ligation of the intersphincteric fistula tract plus a bioprosthetic anal fistula plug procedure in patients with transsphincteric anal fistula: Early results of a multicenter prospective randomized trial. *Annals of Surgery*. 2016;264(6):917–22.
72. Zwiep TM, Gilbert R, Boushey RP, et al. Comparison of ligation of the intersphincteric fistula tract and biolift for the treatment of transsphincteric anal fistula: A retrospective analysis. *Diseases of the Colon and Rectum*. 2020;3:365–70.
73. Madbouly KM, Emile SH, Issa YA, Omar W. Ligation of intersphincteric fistula tract (LIFT) with or without injection of platelet-rich plasma (PRP) in management of high trans-sphincteric fistula-in-ano: Short-term outcomes of a prospective, randomized trial. *Surgery (United States) [Internet]*. 2021;170(1):61–6. Available from: <https://doi.org/10.1016/j.surg.2020.12.025>
74. Narang SK, Keogh K, Alam NN, Pathak S, Daniels IR, Smart NJ. A systematic review of new treatments for cryptoglandular fistula in ano. *Surgeon [Internet]*. 2017;15(1):30–9. Available from: <http://dx.doi.org/10.1016/j.surge.2016.02.002>

75. Jayne DG, Scholefield J, Tolan D, et al. Anal fistula plug versus surgeon's preference for surgery for trans-sphincteric anal fistula: The FIAT RCT. *Health Technology Assessment*. 2019;23(21):1–75.
76. Lin H, Jin Z, Zhu Y, Diao M, Hu W. Anal fistula plug vs rectal advancement flap for the treatment of complex cryptoglandular anal fistulas: a systematic review and meta-analysis of studies with long-term follow-up. *Colorectal Disease*. 2019;21:502–515.
77. de la Portilla F, Dios-Barbeito S, Maestre-Sánchez MV, et al. Feasibility and safety of calcium alginate hydrogel sealant for the treatment of cryptoglandular fistula-in-ano: phase I/IIa clinical trial. *Colorectal Disease*. 2021;23(6):1499–506.
78. Gautier M, Godeberge P, Ganansia R, et al. Easy clip to treat anal fistula tracts: a word of caution. *International Journal of Colorectal Disease*. 2015;30(5):621–4.
79. Marinello F, Kraft M, Ridaura N, Vallribera F, Espín E. Treatment of Fistula-in-ano With OTSC® Proctology Clip Device: Short-term Results. *Cirugia Espanola*. 2018;96(6):369–74.
80. Prosst R. Minimally invasive surgical clip closure of anorectal fistulas: current status of OTSC Proctology. *Minimally Invasive Therapy and Allied Technologies [Internet]*. 2019;28(5):261–7. Available from: <https://doi.org/10.1080/13645706.2018.1521837>
81. Garg P, Garg M. PERFACT procedure: A new concept to treat highly complex anal fistula. *World Journal of Gastroenterology*. 2015;21(13):4020–9.
82. Chase TJG, Quddus A, Selvakumar D, Cunha P, Cuming T. VAAFT for complex anal fistula: a useful tool, however, cure is unlikely. *Techniques in Coloproctology [Internet]*. 2021;25(10):1115–21. Available from: <https://doi.org/10.1007/s10151-021-02492-x>
83. Zhang Y, Li F, Zhao T, Cao F, Zheng Y, Li A. Treatment of Complex Anal Fistula by Video-Assisted Anal Fistula Treatment Combined with Anal Fistula Plug: A Single-Center Study. *Surgical Innovation*. 2021;(16).
84. Zhang Y, Li F, Zhao T, Cao F, Zheng Y, Li A. Video-assisted anal fistula treatment combined with anal fistula plug for treatment of horseshoe anal fistula. *Journal of International Medical Research*. 2021;49(1).
85. Zelić M, Karlović D, Kršul D, Bačić D, Warusavitarne J. Video-Assisted Anal Fistula Treatment for Treatment of Complex Cryptoglandular Anal Fistulas with 2 Years Follow-Up Period: Our Experience. *Journal of Laparoendoscopic and Advanced Surgical Techniques*. 2020;30(12):1329–33.
86. Giarratano G, Shalaby M, Toscana C, Sileri P. Video-assisted anal fistula treatment for complex anal fistula: a long-term follow-up study. *Colorectal Disease*. 2020;22(8):939–44.
87. Regusci L, Fasolini F, Meinero P, et al. Video-Assisted Anal Fistula Treatment (VAAFT) for complex anorectal fistula: efficacy and risk factors for failure at 3-year follow-up. *Techniques in Coloproctology [Internet]*. 2020;24(7):741–6. Available from: <https://doi.org/10.1007/s10151-020-02213-w>
88. la Torre M, Lisi G, D'Agostino E, et al. Lift and VAAFT for high trans-sphincteric anal fistula: a single center retrospective analysis. *International Journal of Colorectal Disease*. 2020;35(6):1149–53.
89. Liu H, Tang X, Chang Y, Li A, Li Z, Xiao Y, et al. Comparison of surgical outcomes between video-assisted anal fistula treatment and fistulotomy plus seton for complex anal fistula: A propensity score matching analysis - Retrospective cohort study. *International Journal of Surgery [Internet]*. 2020;75(January):99–104. Available from: <https://doi.org/10.1016/j.ijssu.2020.01.137>
90. Nordholm-Carstensen A, Perregaard H, Hagen KB, Krarup PM. Fistula Laser Closure (FiLaC™) for fistula-in-ano—yet another technique with 50% healing rates? *International Journal of Colorectal Disease*. 2021;36(9):1831–7.
91. Giamundo P, de Angelis M. Treatment of anal fistula with FiLaC®: results of a 10-year experience with 175 patients. *Techniques in Coloproctology [Internet]*. 2021;25(8):941–8. Available from: <https://doi.org/10.1007/s10151-021-02461-4>
92. Wolicki A, Jäger P, Deska T, et al. Sphincter-saving therapy for fistula-in-ano: long-term follow-up after FiLaC®. *Techniques in Coloproctology [Internet]*. 2021;25(2):177–84. Available from: <https://doi.org/10.1007/s10151-020-02332-4>
93. Brabender DE, Moran KL, Brady Met al. Assessing the effectiveness of laser fistulectomy for anal fistula: a retrospective cohort study. *Techniques in Coloproctology [Internet]*. 2020;24(10):1071–5. Available from: <https://doi.org/10.1007/s10151-020-02281-y>
94. Stijns J, van Loon YT, Clermonts SHEM, et al. Implementation of laser ablation of fistula tract (LAFT) for perianal fistulas: do the results warrant continued application of this technique? *Techniques in Coloproctology*. 2019;23(12):1127–32.
95. Marref I, Spindler L, Aubert M, et al. The optimal indication for FiLaC® is high trans-sphincteric fistula-in-ano: a prospective cohort of 69 consecutive patients. *Techniques in Coloproctology [Internet]*. 2019;23(9):893–7. Available from: <https://doi.org/10.1007/s10151-019-02077-9>
96. de Hous N, de Gheldere C, van den Broeck S, Komen N. FiLaC™ as a last, sphincter-preserving resort for complex perianal fistula. *Techniques in Coloproctology [Internet]*. 2019;23(9):937–8. Available from: <https://doi.org/10.1007/s10151-019-02070-2>
97. Wu YF, Zheng BC, Chen Q, et al. Video-Assisted Modified Ligation of the Intersphincteric Fistula Tract, an Integration of 2 Minimally Invasive Techniques for the Treatment of Parks Type II Anal Fistulas. *Surgical Innovation*. 2021;28(4):419–26.
98. Dong X, Jia Z, Yu B, Zhang X, Xu F, Tan L. Effect of intersphincteric fistula tract ligation versus anal fistulectomy on pain

scores and serum levels of vascular endothelial growth factor and interleukin-2 in patients with simple anal fistulas. *Journal of International Medical Research [Internet]*. 2020 [cited 2021 Sep 1];48(9):1–12.

Available from: <https://us.sagepub.com/en-us/nam/open-access-at-sage>

99. al Sebai OI, Ammar MS, Mohamed SH, el Balshy MA, Ch MBB, el Balshy MA. Comparative study between intersphincteric ligation of perianal fistula versus conventional fistulotomy with or without seton in the treatment of perianal fistula: A prospective randomized controlled trial. *Annals of Medicine and Surgery [Internet]*. 2021 Jan 1 [cited 2021 Oct 11];61(November 2020):180–4. Available from: <https://doi.org/10.1016/j.amsu.2020.12.014>

Table legends

Table 1 – Recent studies assessing VAAFT in anal fistula (number of patients, type of fistula and internal opening closure type)

Table 2 – St James’s University Hospital classification (42)

Table 3 – Recent studies assessing VAAFT in anal fistula (number of procedures and results)

Table 4 – Result of fistula laser closure (FiLaC) in recent studies

Table 5 – Recent randomized controlled trials (RCT) where LIFT procedure was compared or used as comparator.

Table 1

Fistula grade	Description
Grade I	Simple linear intersphincteric fistula
Grade II	Intersphincteric fistulas with abscess or secondary track
Grade III	Trans-sphincteric fistula
Grade IV	Trans-sphincteric fistula with abscess or secondary track within the ischiorectal fossa
Grade V	Supralelevator and translevator fistula

Table 2

Authors	No. patients	Types of anal fistula	Patients with Crohn’s disease	Previous surgery	Closure of the internal opening
Chase et al. (82)	84	complex	11 (13%)	66 (78%)	Vicryl suture or mucosal flap
Zhang et al. (83)	57	complex	-	24 (42%)	Anal fistula plug
Zhang et al. (84)	26	horseshoe	-	14 (54%)	Anal fistula plug
Zelic et al. (85)	73	complex	-	19 (26%)	17 - RAF 16 - mattress suture 14 - LIFT
Giarratano et al. (86)	72	complex	-	18 (25%)	mattress suture
Regusci et al. (87)	104	complex	-	25 (24%)	stapler
La Torre et al. (88)	28	high trans- sphincteric	-	3 (11%)	suture
Liu et al. (89)	64	complex	-	36 (53%)	suture
Jiang et al.	52	complex	1 (2%)	31 (60%)	Suture (2 with stapler)
Total	560		12 (2.1%)	236 (42.1%)	

Table 3

Authors	No. of VAAFT	Repeat VAAFT	Complications Clavien-Dindo grade	Healing	No improvement / worse /recurrence	Follow-up
Chase et al. (82)	105	21	7 (6.7%) grd. II and IIIb	19 (23%) and 34 (38%) symptoms improvement	29 (38%)	6 months
Zhang et al. (83)	57	-	Not reported	51 (89.5%)	6 (10.5%)	6-50 months
Zhang et al. (84)	More than 26	2 patients needed more than 3 procedures	Not reported	23 (88.5%)	3 (11.5%)	6 months
Zelic et al. (85)	83	16	1 (1.4%) - grd. III	62 (85%)	11 (15%)	24 months
Giarratano et al. (86)	77	5	8 (11%) - grd. I	64 (86%)	8 (14%)	12 months
Regusci et al. (87)	120	16	9 (8.6%) - grd. I 4 (3.8%) - grd. II	80 (83.3%)	24 (16.7%)	36 months
La Torre et al. (88)	28	Not stated	4 (14%) - grd. I	24 (86%)	4 (14%)	1-6 months
Liu et al. (89)	64	-	2 (3%)	54 (84%)	10 (16%)	3 months
Jiang et al.	52	8 patients needed more than 3 procedures	25 (29%)	44 (85%)	8 (15%)	9 months
Total	586		60 (10.7%)	455 (81.2%)	103 (18.4%)	

Table 4

Author	No. patients	Crohn's disease patients	Previous surgery before FiLaC	Primary healing rate	Overall healing rate	Follow-up - months	Complications
Nordholm- Carstensen A et al. (90)	66	11 (16%)	28	64.7%	44.1%	26	1 (1.4%)
Giamundo P et al. (91)	175	-	150	117 (66.8%)	129 (73.7%)	4.8-87.6	17
Wolicki A et al. (92)	83	2 (2.4%)	76 (91.6%)	62 (74.7%)	64 (78.3%)	13-84	18
Brabender DE et al. (93)	18	4 (22%)	15 (83.3%)	22%	4 (22%)	18-46	3 (16%)
Serin KR et al. (61)	35	-	8 (38%)	na	15 (42.9%)	6-17.6	na
LaurettaA et al. (62)	30	-	22 (73.3%)	10 (33.3%)	12 (40%)	6-24	4 (13%)
Stijns J et al. (94)	20	-	4 (20%)	4 (20%)	6 (30%)	10	4 (20%)
Marref I et al. (95)	69	6 (8.7%)	37 (54.4%)		31 (45.6%)	4.2-9.3	na
De Hous N et al. (96)	10 (15 fistulas)	2	15 (100%)	10 (67%)	12 (80%)	4-26	na
Wilhelm A et al. (60)	117	13 (11.1%)	113 (96.6%)	75 (69.2%)	103 (88%)	25.4	1 (0.8%)
Total	623 (628)	38 (6%)	468 (74.5%)	324 (52%)	405 (65%)		48 (7.7%)

na - not available

Table 5

Authors	Type of fistula	Trial design	No. of patients	Comparator	Primary healing rate/overall healing rate LIFT	Healing rate comparator	Follow-up months
Wu et al. (97)	Parks type 2	RCT	60 (30/30)	Fistula thread closure	-/29 (97%)	27 (90%)	12
Madbouly et al. (73)	trans-sphincteric fistulas involving >50% of the sphincter	RCT	98 (49/49)	LIFT+ platelet-rich plasma	32/29 (65%/59%)	42/38 (85.7%/77.5%)	12
Dong X et al. (98)	Simple fistulas	RCT	100 (50/50)	fistulectomy	-/48 (95.56%)	42 (82.2%)	na
Han JG et al. (71)	High transsphincteric fistulas (involving >30% of the external sphincter)	RCT	235 (118/117)	LIFT+ bioprosthetic anal fistula plug	98 (83.9%)	109 (94%)	9
Al Sebai et al. (99)	Transsphincteric fistulas	RCT	30 (15/15)	Fistulotomy	12 (80%)	14 (93.3%)	6

na - not available